

The impact of temperature on the bionomics of Aedes (Stegomyia) aegypti, with special reference to the cool geographic range margins

Author(s): Eisen L, Monaghan AJ, Lozano-Fuentes S, Steinhoff DF, Hayden MH, Bieringer

PΕ

Year: 2014

Journal: Journal of Medical Entomology. 51 (3): 496-516

Abstract:

The mosquito Aedes (Stegomyia) aegypti (L.), which occurs widely in the subtropics and tropics, is the primary urban vector of dengue and yellow fever viruses, and an important vector of chikungunya virus. There is substantial interest in how climate change may impact the bionomics and pathogen transmission potential of this mosquito. This Forum article focuses specifically on the effects of temperature on the bionomics of Ae. aegypti, with special emphasis on the cool geographic range margins where future rising temperatures could facilitate population growth. Key aims are to: 1) broadly define intra-annual (seasonal) patterns of occurrence and abundance of Ae. aegypti, and their relation to climate conditions; 2) synthesize the existing quantitative knowledge of how temperature impacts the bionomics of different life stages of Ae. aegypti; 3) better define the temperature ranges for which existing population dynamics models for Ae. aegypti are likely to produce robust predictions; 4) explore potential impacts of climate warming on human risk for exposure to Ae. aegypti at its cool range margins; and 5) identify knowledge or data gaps that hinder our ability to predict risk of human exposure to Ae. aegypti at the cool margins of its geographic range now and in the future. We first outline basic scenarios for intra-annual occurrence and abundance patterns for Ae. aegypti, and then show that these scenarios segregate with regard to climate conditions in selected cities where they occur. We then review how near-constant and intentionally fluctuating temperatures impact development times and survival of eggs and immatures. A subset of data, generated in controlled experimental studies, from the published literature is used to plot development rates and survival of eggs, larvae, and pupae in relation to water temperature. The general shape of the relationship between water temperature and development rate is similar for eggs, larvae, and pupae. Once the lower developmental zero temperature (10-14 degrees C) is exceeded, there is a near-linear relationship up to 30 degrees C. Above this temperature, the development rate is relatively stable or even decreases slightly before falling dramatically near the upper developmental zero temperature, which occurs at -38-42 degrees C. Based on life stage-specific linear relationships between water temperature and development rate in the 15-28 degrees C range, the lower developmental zero temperature is estimated to be 14.0 degrees C for eggs, 11.8 degrees C for larvae, and 10.3 degrees C for pupae. We further conclude that available population dynamics models for Ae. aegypti, such as CIMSiM and Skeeter Buster, likely produce robust predictions based on water temperatures in the 16-35 degrees C range, which includes the geographic areas where Ae. aegypti and its associated pathogens present the greatest threat to human health, but that they may be less reliable in cool range margins where water temperatures regularly fall below 15 degrees C. Finally, we identify knowledge or data gaps that hinder our ability to predict risk of human exposure to Ae. aegypti at the cool margins of its range, now and in the future, based on impacts on mosquito population dynamics of temperature and other important factors, such as water nutrient content, larval density, presence of

Climate Change and Human Health Literature Portal

biological competitors, and human behavior.

Source: http://www.ncbi.nlm.nih.gov/pubmed/24897844

Resource Description

Exposure:

weather or climate related pathway by which climate change affects health

Ecosystem Changes, Food/Water Quality, Precipitation, Temperature

Food/Water Quality: Other Water Quality Issue

Water Quality (other): Water temperature

Temperature: Fluctuations

Geographic Feature: M

resource focuses on specific type of geography

General Geographical Feature

Geographic Location:

resource focuses on specific location

Global or Unspecified

Health Impact: M

specification of health effect or disease related to climate change exposure

Infectious Disease

Infectious Disease: Vectorborne Disease

Vectorborne Disease: Mosquito-borne Disease

Mosquito-borne Disease: Chikungunya, Dengue, Yellow Fever

Resource Type: M

format or standard characteristic of resource

Review

Timescale: M

time period studied

Time Scale Unspecified